



**UNIVERSITI PUTRA MALAYSIA**

**PREVALENCE AND MOLECULAR CHARACTERIZATION  
OF VANCOMYCIN-RESISTANT ENTEROCOCCI ISOLATED  
FROM POULTRY, RAW VEGETABLES AND CLINICAL SOURCES**

**OOI WAI LING**

**FSMB 2003 27**

**PREVALENCE AND MOLECULAR CHARACTERIZATION  
OF VANCOMYCIN-RESISTANT ENTEROCOCCI ISOLATED  
FROM POULTRY, RAW VEGETABLES AND CLINICAL SOURCES**

**OOI WAI LING**

**DOCTOR OF PHILOSOPHY  
UNIVERSITI PUTRA MALAYSIA**

**2003**



**PREVALENCE AND MOLECULAR CHARACTERIZATION  
OF VANCOMYCIN-RESISTANT ENTEROCOCCI ISOLATED  
FROM POULTRY, RAW VEGETABLES AND CLINICAL SOURCES**

**By  
OOI WAI LING**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

**August 2003**



## **DEDICATIONS**

**To my beloved parents, brothers, sisters, relatives and friends. Also to those who have taught me and still continue in teaching me of different angles of life. Also to those who have sincerely provided invaluable assistance, guidance, advice, moral support and encouragement through out the whole project including the writing out part.**

**OOI WAI LING 2003**

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment  
of the requirements for the degree of Doctor of Philosophy

**PREVALENCE AND MOLECULAR CHARACTERIZATION  
OF VANCOMYCIN-RESISTANT ENTEROCOCCI ISOLATED  
FROM POULTRY, RAW VEGETABLES AND CLINICAL SOURCES**

By

**OOI WAILING**

**August 2003**

**Chairman: Associate Professor Son Radu, Ph.D.**

**Faculty: Food Science and Biotechnology**

In this study, vancomycin-resistant *Enterococcus* (VRE) were isolated from 120, 60, 180, 450, 850, and 850 with chicken samples (CS), whole chicken (W), hospital clinical samples (HCS), environmental chicken faecal samples (CFS), ulam (U) and vegetables samples, respectively. All these samples were obtained mainly from wet markets and areas around Sri Serdang, Seri Kembangan and Kuala Lumpur. Vancomycin-resistant enterococci isolated from these samples were characterized phenotypically and genotypically. The prevalence of VRE isolated are as follow: HCS (14/180, 7.8%) with 1 isolate as *E. faecalis* (1/14, 7.1%) and 13 as *E. faecium* (13/14, 92.9%); CFS (27/450, 6%) with 24 isolates as *E. raffinosus* (24/27, 88.9%) and 3 as *E. faecium* (3/27, 11.1%); W (8/60, 13.3%) with 1 isolate as *E. faecalis* (1/8, 12.5%), 2 as *E. gallinarum* (3/8, 37.5%), and 4 as *E. faecium* (4/8, 50.0%); CS (16/120, 13.3%) with 2 isolates identified as *E. gallinarum* (2/16, 12.5%), 3 as *E. faecalis* (3/16, 18.8%) and 11 as *E. raffinosus* (11/16, 84.6%); U

(8/850, 0.9%) with 3 isolates as *E. faecium* (3/8, 37.5%) and 5 as *E. raffinosus* (5/8, 62.5%); V (8/850, 0.9%) with 5 as *E. faecium* (5/8, 62.5%) and 3 as *E. gallinarum* (3/8, 37.5%). The MAR index for VRE isolates examined ranged from 0.07-0.93 with *E. raffinosus* having the highest ARI value. Among the 14 antibiotics tested, the highest prevalence of resistance was against kanamycin, streptomycin, erythromycin, bacitracin, tetracycline, and vancomycin (100%); cefuroxime and chloramphenicol (93%), penicillin G (89%), gentamicin (52%), while the lowest resistance was observed for ampicillin (40%), ceftriaxone (34%), rifampicin (10%) and teicoplanin (0%). The VRE isolates generated various plasmid profiles with sizes ranging from 1.7-35.8 MDa. Most isolates were found devoid of plasmid. The multiplex PCR (M-PCR) displayed 3 identified isolates harbouring the *vanA* gene (3.7%), whereas, in single PCR, 42 isolates harboured the *vanA* gene (51.85%). Both M-PCR and single PCR identified 1 isolate (V13) harbouring *vanB* gene, whilst, 16 VRE isolates isolated from CFS, V and U samples were positive for *vanC1* gene. No *vanC2,3*, *vanD*, *vanE* and *vanG* genes were detected among the isolates. Three random primers, GEN 1-50-03 (5'-CTT GAG TGG A-3'), GEN 1-50-05 (5'-TCC TCAAGA C-3'), and GEN 1-50-08 (5'-GAG ATG ACG A-3') used in RAPD-PCR analysis produced bands with molecular size ranging from 0.25 kb to 3.0 kb. PFGE analysis using *SmaI* (5'-CCC↓GGG-3') produced high discrimination fragment patterns among all the VRE isolates examined. Based on the data obtained from the fragment profiles in both RAPD-PCR and PFGE analysis, relationship among the isolates was determined by phylogenetic software (RAPDistance v.1.04). In the immunological evaluation, passive haemagglutination assay (PHA) was used to check immunogenicity. BM4147 gave the highest PHA value followed by CFS66, V583, and BM4174 in the 5 VRE cells treatment groups. The antibody titres in PHA,

demonstrated a relative correlation between the immunogenicity and the VRE cells antigenic properties on the cell walls. The application of PCR (*van* genes) and immunological evaluation methods enable the retrieval of information on both pathogenicity and immunogenicity of VRE isolates examined. Our results in the study demonstrates that the plasmid profiling, M-PCR, and single PCR techniques are able to act as useful analysis tools for rapid and reliable typing and identification of VRE; whilst, the RAPD and PFGE analysis are useful discrimination tool to differentiate the VRE isolates examined.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PREVALEN DAN PENCIRIAN SECARA MOLEKULAR ENTEROCOCCI  
KERINTANGAN VANKOMISIN YANG DIPENCILKAN DARI SUMBER-  
SUMBER TERNAKAN AYAM, SAYURAN MENTAH DAN KLINIKAL**

Oleh

**OOI WAI LING**

**Ogos 2003**

**Pengerusi: Profesor Madya Son Radu, Ph.D.**

**Fakulti: Sains Makanan dan Bioteknologi**

Di dalam kajian ini, *Enterococcus kerintangan* vankomisin (VRE) telah dipencilkan daripada 120, 60, 180, 450, 850, dan 850 dengan sampel ayam kisar (CS), sample ayam basuhan (W), klinikal (HCS), sample persekitaran dari tinja ayam (CFS), ulam (U) dan sampel sayuran. Semua sample ini telah diperolehi daripada pasar-pasar dan kawasan di sekitar Sri Serdang, Seri Kembangan dan Kuala Lumpur. Enterococci kerintangan vankomisin yang telah dipencilkan dicirikan selanjutnya secara fenotipik dan genotipik. Prevalen VRE yang dipencilkan adalah seperti berikut: HCS (14/180, 7.8%) dengan satu isolate sebagai *E. faecalis* (1/14, 7.1%), manakala tiga belas sebagai *E. faecium* (13/14, 92.9%); CFS (27/450, 6%), 24 isolate dikenalpasti sebagai *E. raffinosus* (24/27, 88.9%) dan tiga sebagai *E. faecium* (3/27, 11.1%); W (8/60, 13.3%) dengan satu isolate dikenalpasti sebagai *E. faecalis* (1/8, 12.5%), 2 sebagai *E. gallinarum* (3/8, 37.5%), dan 4 sebagai *E. faecium* (4/8, 50.0%); CS (16/120, 13.3%) dengan 2 sebagai *E. gallinarum* (2/16, 12.5%), 3 sebagai *E. faecalis* (3/16, 18.8%) dan 11 sebagai *E. raffinosus* (11/16, 84.6%); U (8/850, 0.9%) dengan 3 isolate dikenalpasti sebagai *E. faecium* (3/8, 37.5%) dan 5



sebagai *E. raffinosus* (5/8, 62.5%); V (8/850, 0.9%), dengan 5 dikenalpasti sebagai *E. faecium* (5/8, 62.5%) dan 3 sebagai *E. gallinarum* (3/8, 37.5%). Index MAR bagi isolate VRE mempunyai nilai julat 0.07-0.93 manakala *E. raffinosus* adalah spesies yang memberikan nilai ARI tertinggi. Daripada 14 antibiotik yang dikaji, prevalen kerintangan tertinggi dalam isolate VRE kajian adalah terhadap kanamisin, streptomisin, erithromisin, basitrasin, tetrasiklin dan vankomisin (100%); sefuroxime dan kloramfenikol (93%), penisillin G (89%), gentamisin (52%), manakala kerintangan terendah untuk ampicillin (40%), seftriaxon (34%), rifampisin (10%) dan teikoplanin (0%). Isolate VRE memberikan profil plasmid dengan julat saiz dari 1.7-35.8 MDa. Kebanyakan isolate menunjukkan ketidakhadiran plasmid. Analisis multiplex PCR (M-PCR) memaparkan tiga isolate yang dikenalpasti menunjukkan kehadiran gen *vanA* (3.7%) berbanding 42 isolate yang positif untuk gen *vanA* dalam PCR tunggal (51.85%). Kedua-dua analisis M-PCR dan PCR tunggal mengenalpasti 1 isolate yang positif kepada gen *vanB* (V13), manakala 16 isolate daripada CFS, V dan U adalah positif kepada gen *vanC1*. Tiada sebarang gen yang dikesan untuk gen *vanC2,3*, *vanD*, *vanE* dan *vanG* untuk kesemua isolate yang dikaji. Tiga primer rawak, GEN 1-50-03 (5'-CTT GAG TGG A-3'), GEN 1-50-05 (5'-TCC TCAAGA C-3'), dan GEN 1-50-08 (5'-GAG ATG ACG A-3') yang digunakan dalam analisis RAPD menghasilkan fragmen yang bersaiz molekul antara 0.25 kb-3.0 kb. Analisis yang menggunakan enzim pemotong *SmaI* (5'-CCC↓GGG-3'), menghasilkan fragmen corak yang berdiskriminasi tinggi di antara semua isolate VRE yang dikaji. Berdasarkan maklumat yang diperolehi daripada profil fragmen di dalam kedua-dua analisis RAPD-PCR dan PFGE, hubungan di antara isolate telah dikenalpastikan dengan menggunakan cakera ringan untuk analisis filogenetik (RAPDistance v.1.04) Di dalam Kajian immunology, kaedah hemagglutinasin pasif (PHA) telah digunakan

untuk mengkaji immunogenisiti. BM4147 memberikan nilai PHA yang tertinggi berikutan dengan CFS66, V583, dan BM4174 di dalam lima kumpulan rawatan sel VRE. Titer antibodi memaparkan korelasi relatif di antara immunogenisiti dan kesan antigenik pada permukaan dinding sel VRE. Aplikasi kaedah PCR (gen *van*) dan kaedah kajian immunology membolehkan maklumat tentang patogenisiti dan imunogenisiti diperolehi untuk kajian isolate VRE. Keputusan kita turut menunjukkan kaedah profil plasmid, M-PCR, dan PCR tunggal adalah berkemampuan digunakan sebagai alat penganalisaan yang cepat, berkesan dan bersesuaian bagi pencirian dan pengenalan VRE; manakala analisis dengan RAPD-PCR dan PFGE amat berguna sebagai alat perbezaan untuk membezakan isolate VRE yang dikaji.

## ACKNOWLEDGEMENTS

I would particularly like to express my deepest gratitude to my respectable supervisor, Associate Professor Dr. Son Radu for his continuous strong support, patience, invaluable advice, and guidance throughout the course of this project. My deepest gratitude goes to my two other supervisory committees, Professor Dr. Gulam Rusul Rahmat Ali and Professor Dato' Dr. Sheikh Omar Abdul Rahman. Without their persistent assistance and exceptional generosity, this work would not have been possible and meet the requirements of this course.

Special appreciation goes to all the postgraduate students from the Food Safety and Molecular Subtyping Group for their continuous support and their generosity to help out throughout the whole project.

Heartfelt thanks to those helpful staffs and students from the Department of Plant Protection, Faculty of Agriculture; also from the Malaysian Agricultural Research and Development Institute (MARDI) for the generosity to share the information and laboratory facilities for bacterial identification purposes. Also, not forgetting to those who had contributed either directly or indirectly in the success of my project work, thanks a lot!

Last but not least, I would like to extend my deepest gratitude to the most supportive group of my truly loving family members and all my friends for their invaluable supports, encouragements and sharing of all my joys and sorrows in life. This did indeed sustained and motivated me not only for this period but will continuously inspired me in future.

Finally, words just can't express what is beholding in my heart, please accept my sincere thankfulness and gratitude for everything.

I certify that an Examination Committee met on 29<sup>th</sup> August, 2003 to conduct the final examination of Ooi Wai Ling on her Doctor of Philosophy thesis entitled “Prevalence and Molecular Characterization of Vancomycin-Resistant Enterococci Isolated from Poultry, Raw Vegetables and Clinical Sources” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committees are as follows:

**Raha Abdul Rahim, Ph.D.**

Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Chairman)

**Son Radu, Ph.D.**

Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Main Supervisor)

**Gulam Rusul Rahmat Ali, Ph.D.**

Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**Dato’ Sheikh Omar Abdul Rahman, Ph.D.**

Professor  
Faculty Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

**Giorgio Giraffa, Ph.D.**

Professor  
Department Microbiology and Enzymology  
Institute of Experimental Lattiero Caseario  
Italy  
(External Examiner)



**GULAM RUSUL RAHMAT ALI, Ph.D.**  
Professor / Deputy Dean  
School of Graduate Studies,  
Universiti Putra Malaysia.

Date: 30 SEP 2003

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

**Son Radu, Ph.D.**

Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Chairman)

**Gulam Rusul Rahmat Ali, Ph.D.**

Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**Dato' Sheikh Omar Abdul Rahman, Ph.D.**

Professor  
Faculty Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

**Giorgio Giraffa, Ph.D.**

Professor  
Department Microbiology and Enzymology  
Institute of Experimental Lattiero Caseario  
Italy  
(External Examiner)



---

**AINI IDERIS, Ph.D**  
Professor / Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: **14 NOV 2003**

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

  
(OOI WAI LING)

Date: 26/10/2003

## TABLE OF CONTENTS

	<b>Page</b>
DEDICATIONS	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL SHEETS	x
DECLARATION FORM	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xviii
LIST OF ABBREVIATIONS	xx
 <b>CHAPTER</b>	
 <b>I      GENERAL INTRODUCTION</b>	 <b>1</b>
 <b>II     LITERATURE REVIEW</b>	
Taxonomy and Classification	7
The Genus <i>Enterococcus</i>	8
Serological Identification	10
Natural Habitat	12
Pathogenicity	13
Identification	14
Vancomycin-Resistant Enterococci	17
Epidemiology and Transmission of VRE	20
Phenotyping Techniques	
Antibiotic Susceptibility Testing / Disk Diffusion Testing	21
E-test Strips Testing	24
Plasmid Profiling	
Plasmid	24
Molecular Genotyping	
Polymerase Chain Reaction	26
Multiplex PCR	29
Vancomycin-Resistant Genes ( <i>vanA</i> , <i>vanB</i> , <i>vanC1</i> , <i>vanC2,3</i> , <i>vanD</i> , <i>vanE</i> , <i>vanG</i> )	30
RAPD Fingerprinting	31
Agarose Gel Electrophoresis	34
Macrorestriction Analysis of DNA by Pulsed-Field Gel Electrophoresis	35
Genetic Statistical Analysis	37
Immunological Evaluation Application	
Animal Preparation (Selection & Injection)	38
Sample Collection	39

	Sample Collection	39
	Sensitisation of Sheep Red Blood Cells	39
	PHA Testing	39
<b>III</b>	<b>ISOLATION AND IDENTIFICATION OF VANCOMYCIN RESISTANT ENTEROCOCCI</b>	
	Introduction	40
	Materials and Methods	42
	Results	44
	Discussion	47
<b>IV</b>	<b>RAPID DETECTION OF VANCOMYCIN GENOTYPES OF <i>vanA</i>, <i>vanB</i>, <i>vanC1</i>, <i>vanC2,3</i>, <i>vanD</i>, <i>vanE</i> and <i>vanG</i> UTILISING MODIFIED MULTIPLEX PCR (M-PCR) AND SINGLE PCR METHOD AS A COMPARISON</b>	
	Introduction	51
	Materials and Methods	
	PCR Amplification Procedures	53
	Agarose Gel Electrophoresis	56
	Results	57
	Discussion	64
<b>V</b>	<b>ANTIBIOTICS SUSCEPTIBILITY TESTING AND PLASMID PROFILING</b>	
	Introduction	69
	Materials and Methods	
	Antibiotic Susceptibility Testing (Disk Diffusion Test)	72
	Antibiotic Sensitivity Analysis	73
	Antibiotic Resistance Indexing (ARI) of VRE Species	73
	Multiple Antibiotic Resistance (MAR) Indexing of VRE isolates	73
	E-Testing	74
	Plasmid Analysis	74
	Results	76
	Discussion	87
<b>VI</b>	<b>RANDOM AMPLIFIED POLYMORPHIC DNA (RAPD) FINGERPRINTING</b>	
	Introduction	91
	Materials and Methods	
	RAPD-PCR Amplification	92
	Agarose Gel Electrophoresis and UV Photographing	94
	Fingerprinting pattern Analysis	95
	Results	96
	Discussion	103



<b>VII</b>	<b>PULSED-FIELD GEL ELECTROPHORESIS (PFGE)</b>	
	Introduction	107
	Materials and Methods	109
	Rapid DNA Plug Preparation	110
	Restriction Enzyme Digestion	110
	PFGE Agarose Gel Electrophoresis	111
	Data Analysis	111
	Results	112
	Discussion	116
<b>VIII</b>	<b>IMMUNOLOGICAL EVALUATION</b>	
	Introduction	120
	Materials and Methods	
	Animal Preparation	122
	Sample Collection	122
	Sensitisation of sRBC	123
	PHA Testing	124
	Results	125
	Discussion	125
<b>IX</b>	<b>GENERAL DISCUSSION</b>	128
<b>X</b>	<b>CONCLUSION</b>	133
	<b>REFERENCES</b>	135
	<b>APPENDICES</b>	
	A. Culture Media and Reagents Preparation	160
	B. Figures of results	165
	<b>VITAE</b>	

## LIST OF TABLES

Table	Page
2.1 Members of <i>Enterococcus</i> groups	11
2.2 The identification chart for <i>Enterococcus</i> groups (adopted from Facklam and Collins, 1989)	15
2.3 Chart showing the identification and biochemical reaction response of enterococcal species in conventional testing	16
3.1 The prevalence of VRE isolates.	45
3.2 The prevalence of VRE species obtained in the study.	46
3.3 The distribution of VRE positive species in different sample sources	47
4.1 Primer pairs utilized in the multiplex PCR. Set 1 M-PCR mixture: <i>vanA</i> , <i>vanB</i> and <i>vanC1</i> ; Set 2 M-PCR mixture: <i>vanC2/C3</i> , <i>vanD</i> , <i>vanE</i> and <i>vanG</i> . (S, sense; AS, antisense)	55
4.2 The distribution for all the positive vancomycin genes obtained over the total VRE isolates studied for both total sample population and total VRE isolates. There are no <i>vanC23</i> , <i>vanD</i> , <i>vanE</i> and <i>vanG</i> genes detected from all the sample isolates examined. TSI, total sample population; TVI, total VRE isolates.	59
5.1 Groups of antimicrobial used in the study and their respective mechanism of actions	70
5.2 The distribution of antimicrobial resistance of all the isolates of VRE used in the study	78
5.3 The antibiotic resistance, plasmid profiling and MIC value (E-test) for all the VRE isolates (CFS, HCS, CS, W, U, V) in the study including both the positive and negative control VRE cultures (V583, BM4147, BM4174, and ATCC 19434)	82
5.4(a) The antibiotic resistance profile patterns obtained can be grouped into types as observed above. The MAR index value was also included	83
5.4(b) The antibiotic resistance profile patterns grouped from A-K with the respective VRE isolates	84
5.5 Antibiotic resistance among VRE species isolates to 14 antibiotics used in the study	85

5.6	The plasmid profile patterns. The size reference marker for plasmid (produced by <i>E. coli</i> V517: 35.8, 4.8, 3.7, 3.4, 2.6, 2.0, 1.8, and 1.4 MDa)	85
5.7	The VRE strains used in the study for antibiogram and plasmid profiling	86
6.1	Random primers used in the random amplification polymorphic DNA (RAPD) procedure ( <i>GENOSYS</i> , Biotechnologies Inc, USA). Only three primers GEN1-50-03, GEN1-50-05 and GEN1-50-08 were chosen for further characterization study for all VRE isolates in this study as they yielded clear and highly polymorphic banding profiling	94
8.1	The antibody titres for each test serum from week 0 to week 8 in 10 rabbits.	127

## LIST OF FIGURES

Figure	Page
2.1 Phylogenetic relationship between varying genera similar to <i>Enterococcus</i> genus (adopted from Scheifer and Ludwig, 1995)	9
2.2 Identification chart for vancomycin-resistant <i>Enterococcus</i> species using conventional tests	18
2.3 The chemical structure of vancomycin	19
2.4 The basic PCR procedures involving three major steps. (source: PCR technical bulletin, Promega corporation)	28
2.5 Arbitrary primed PCR is applied in the generation of polymorphic fingerprints for the genetic mapping. At low stringency, oligonucleotide primers are annealed to the template. This reaction depends on the DNA sequence of the template and the sequence polymorphisms are influencing the details of the interaction occurs	33
4.1 Electrophoresis profiles obtained for the M-PCR assaying. Lane M, 100bp DNA ladder, lanes 1-16: VRE isolates for U106, VE13, V48, CFS 23, CFS29, CFS 30, CFS 35, CFS 39, CFS 47, CFS 49, CFS 61, CFS 65, BM4174 ( <i>vanC</i> , positive control), V583 ( <i>vanB</i> , positive control), <i>E. faecalis</i> ( <i>vanB</i> , normal control), ATCC 19434	61
4.2 Electrophoresis profiles obtained for the M-PCR assaying, lane M, 100bp DNA ladder; lanes 1-8: CFS66, CFS122, CFS123, V13, BM4147 ( <i>vanA</i> , positive control), clinical known <i>E. faecium</i> ( <i>vanA</i> , Normal control), HCS22, U106	61
4.3 Electrophoresis profiles obtained for the single PCR assaying for <i>vanA</i> . Lane M, 100bp DNA ladder; lanes 1-25: VRE isolates for CFS23, CFS29, CFS30, CFS35, CFS39, CFS42, CFS47, CFS49, CFS50, CFS51, CFS59, CFS61, CFS65, CFS66, CFS122, CFS123, HCS1, HCS4, HCS6, HCS7, HCS10, HCS11, PC (BM4147), NC (ATCC19434)	62
4.4 Electrophoresis profiles obtained for the single PCR assaying for <i>vanB</i> . Lane M, 100bp DNA ladder; lanes 1-21: VRE isolates for U106, U107, U112, U114, U133, U134, U157, U158, V13, V4b, V27, V39, V40, V48, V50, V55, W7, W8, W14, PC (BM4174), NC (ATCC19434)	62
4.5 Electrophoresis profiles obtained for the single PCR assaying for <i>vanC1</i> . Lane M, 100bp DNA ladder, lanes 1-17: VRE isolates for CFS23, CFS29, CFS30, CFS35, CFS39, CFS42, CFS47, CFS49,	

	CFS50, CFS51, CFS59, CFS61, CFS65, V13, V48, U106, NC (ATCC19434)	63
5.1	The gel electrophoresis of plasmid profiling for the first 8 VRE isolates Of chicken faecal samples. Lane M, plasmid reference marker, <i>E. coli</i> V517. Lanes 1-14: CFS23, CFS29, CFS30, CFS35, CFS39, CFS42, CFS47, CFS49, CFS 50, CFS59, CFS60, CFS61, CFS65, ATCC 19434	79
6.1	Gel shows the RAPD fingerprinting profile obtained for VRE isolates of CFS samples utilizing GEN 1-50-08 and GEN 1-50-03. Lane M, 1 kb DNA ladder marker; lanes 1-16: CFS23, CFS29, CFS30, CFS35, CFS39, CFS42, CFS47, CFS49 (primer GEN 1-50-08); CFS50, CFS59, CFS60, CFS61, CFS65, CFS66, CFS122, CFS123 (primer GEN 1-50-03)	97
6.2	Gel shows the RAPD fingerprinting profile obtained for VRE isolates of HCS samples utilizing GEN 1-50-03 and GEN 1-50-05. Lane M, 1 kb DNA ladder marker; lanes 1-16: HCS1, HCS4, HCS6, HCS7, HCS10, HCS11, HCS14 (primer GEN 1-50-03); HCS1, HCS4, HCS6, HCS7, HCS10, HCS11, HCS14 (primer GEN 1-50-05)	97
6.3	Dendrogram showing the genetic relatedness among all the 81 VRE isolates obtained from six different sources of food, clinical and environmental samples	98
6.4	The dendrogram showing the genetic relatedness among all the VRE isolates of <i>E. raffinosus</i> obtained from different sample sources	99
6.5	The dendrogram showing the genetic relatedness among all the VRE isolates of <i>E. faecium</i> obtained from different sample sources	100
6.6	The dendrogram showing the genetic relatedness among all the VRE isolates of <i>E. faecalis</i> obtained from different sample sources	101
6.7	The dendrogram showing the genetic relatedness among all the VRE isolates of <i>E. gallinarum</i> obtained from different sample sources	102
7.1	Gel shows the PFGE analysis of VRE isolates from chicken faecal samples (CFS). Lane 1 to 14: CFS23, CFS29, CFS30, CFS35, CFS39, CFS42, CFS47, CFS49, CFS51, CFS59, CFS61, CFS65, CFS122, and CFS123; lane M: PFGE molecular size marker	114
7.2	Gel shows the PFGE analysis of VRE isolates from hospital clinical samples (HCS). Lane 1 to 14: HCS1, HCS4, HCS6, HCS7, HCS9, HCS10, HCS11, HCS14, HCS15, HCS17, HCS18, HCS20, HCS21, HCS22; lane M: PFGE molecular size marker	114
7.1	The dendrogram generated from PFGE profiles showing the genetic relatedness among all the VRE isolates from six different sources of food, clinical and environmental samples	115

## LIST OF ABBREVIATIONS

### Abbreviations

A	adenine or adenosine
AP-PCR	arbitrarily primed-polymerase chain reaction
ATCC	American Type Culture Collection
ATP	adenosine triphosphate
Amp	ampicillin
B	bacitracin
bp	basepair
BSA	bovine Serum Albumin
C	chloramphenicol
ccc	covalently closed circular
cm	centimetre
Cro	ceftriaxone
Cxm	cefuroxime
Da	dalton
dATP	deoxyadenosine triphosphate
dCTP	deoxycytosine triphosphate
dGTP	deoxyguanine triphosphate
dTTP	deoxythymidine triphosphate
dH <sub>2</sub> O	distilled water
DNA	deoxyribonucleic acid
dNTPs	deoxy nucleotide triphosphate (PCR nucleotide mix containing dATP, dTTP, dGTP and dCTP)
E	erythromycin

<i>E.coli</i>	<i>Escherichia coli</i>
e.g.	for example
EDTA	ethylenediamine tetraacetic acid
EtBr	ethidium bromide
g	gram
g	gravity
G	guanine
Gm	gentamicin
h	hour
i.e.	that is
I	intermediate
ID	identification number
K	kanamycin
kb	kilobase
KDa	kiloDalton
kg	kilogram
LB	Luria-Bertani
M	molar, or molarity, moles of solute per litre of solution
MDa	megaDalton
mg	milligram
MHA	Mueller Hinton Agar
min	minutes
ml	millilitre
mm	millimetre
mM	miliMolar

µg	microgram
µl	microlitre
mol	mole
Na	sodium
NaCl	sodium chloride
NaOH	sodium hydroxide
P	penicillin
%	percentage
PCR	polymerase chain reaction
PFGE	pulsed-field gel electrophoresis
R	resistant
RAPD	random amplified polymorphic DNA
Rd	rifampicin
RE	restriction enzyme
REA	restriction enzyme analysis
RFLP	restriction fragment length polymorphism
RNA	ribonucleic acid
RNase	ribonuclease
rpm	rotation per minute
s	sensitive
S	streptomycin
sdH <sub>2</sub> O	sterile distilled water
SDS	sodium dodecyl sulphate
<i>Taq</i>	<i>Thermus aquaticus</i> DNA (polymerase)
TBE	tris borate EDTA electrophoresis buffer



<b>TE</b>	<b>tris EDTA buffer</b>
<b>Te</b>	<b>tetracycline</b>
<b>Tec</b>	<b>teicoplanin</b>
<b>Tris</b>	<b>tris (hydroxymethyl) methylamine</b>
<b>UV</b>	<b>ultraviolet</b>
<b>V</b>	<b>volts</b>
<b>Va</b>	<b>vancomycin</b>
<b>v/v</b>	<b>volume per volume</b>
<b>w/v</b>	<b>weight per volume</b>